

ENVIRONMENTAL CARCINOGENESIS

Lifestyle Factors, Exposures, Genetic Susceptibility, and Renal Cell Cancer Risk: A Review

Lee E. Moore and Robin T. Wilson

*Occupational Epidemiology Branch, Division of Cancer Epidemiology and Genetics,
National Cancer Institute, NIH, DHHS, Bethesda, Maryland, USA*

Sharan L. Campleman

Public Health Institute, California Cancer Registry, Sacramento, California, USA

Malignant kidney tumors account for approximately 2% of all new primary cancer cases diagnosed in the United States, with an estimated 30,000 cases occurring annually. Although a variety of agents, chemical and biological, have been implicated as causal agents in the development of renal cell carcinoma (RCC), the etiology remains enigmatic. The strongest association has been developed between cigarette smoking and renal cancer however consistent, positive associations between RCC and obesity, diabetes, and hypertension have also been reported. In addition, more recent investigations of familial kidney cancer syndromes indicate that a strong genetic component contributes to RCC development. Several genes have been identified through investigation of familial kidney cancer syndromes. This review article describes recent trends in RCC incidence and the currently identifiable etiological causes that account for approximately half of the RCC cases diagnoses. The remainder of this review then focuses on additional risk factors that have thus far not been well examined but may be helpful in explaining the increasing incidence trends and the geographic or racial variation observed nationally and worldwide.

Keywords Kidney cancer; Exposures; Genetic susceptibility; Review

INTRODUCTION

Malignant tumors of the kidney account for approximately 2% of all new primary cancer cases diagnosed in the United States, with an estimated 30,000 cases occurring annually.^[1,2] Renal cell carcinoma (RCC) of the renal parenchyma accounts for >80% of all renal cancers, the majority of which are adenocarcinomas.^[3] RCC is subdivided into several histologic subtypes. The largest group, clear cell carcinoma, accounts for 80–85% of cases followed by papillary carcinoma, accounting

for about 10%. The remaining types, such as oncocytomas and chromophobe tumors, are more rare. Other less common histological types include transitional cell malignancies, most often occurring in the renal pelvis; nephroblastoma or Wilms' tumor, an embryonal malignancy of early childhood; and a mixture of sarcomas.^[3]

Although a variety of agents, chemical and biological, have been implicated as causal agents in the development of RCC, the etiology, nonetheless, remains enigmatic. The strongest association has been developed between cigarette smoking and renal cancer, for both cancers of the renal parenchyma (usually referred to as renal cell) and those of the renal pelvis and ureter.^[4,5] Consistent, positive associations between RCC and obesity (high relative weight), diabetes, and hypertension have also been reported.^[6,7] Other risk factors associated with renal cancer include abuse of analgesics,^[4,8] history of cystic disease due to long-term hemodialysis,^[9] and certain occupations.^[10,11]

In addition, more recent investigations of familial kidney cancer syndromes indicate that a strong genetic component contributes to RCC development.^[12,13] Several genes have been identified through investigation of familial kidney cancer syndromes. The relationship between the von Hippel-Lindau (VHL) tumor suppressor gene and clear cell carcinoma,^[12] the MET proto-oncogene gene and papillary carcinoma,^[13] and other genes associated with inherited syndromes, such as familial oncocytoma^[14] and the Birt-Hogg-Dubé syndrome^[15] have provided new insights into the causal mechanism of each disease and may also be important in understanding sporadic forms of the disease.

In combination with the enigmatic nature of RCC etiology, several factors lead to a high potential for substantial cancer-related health disparities, including the relatively high occurrence of RCC among certain minorities, the large racial and geographic variations in disease incidence, the underlying genetic susceptibilities, and the continued high proportion of cases diagnosed at advanced stage, a factor related to low survival. This review article briefly describes recent trends in

Address correspondence to Lee E. Moore, Occupational Epidemiology Branch, Division of Cancer Epidemiology and Genetics, National Cancer Institute, NIH, DHHS, Bethesda, MD, USA; Fax: (301) 402-1819; E-mail: moorele@mail.nih.gov




FIG. 1. Age-adjusted (2000 U.S. Standard) incidence rates for renal cell carcinoma per 100,000 person-years by sex and race/ethnicity in the United States—NCI-SEER, 1974–1999 and CCR, 1992–1999.

RCC incidence and the currently identifiable etiological causes that account for approximately half of the RCC case diagnoses. The remainder of this review then focuses on additional risk factors that have thus far not been well examined but may be helpful in explaining the increasing incidence trends and the geographic or racial variation observed nationally and worldwide.

INCIDENCE, TRENDS, AND RACIAL VARIATION

Recent descriptive studies have reported increases in the overall incidence of kidney cancer since the early 1970s, not only in the United States but also globally including portions of Asia, Oceania, Europe, and South America.^[16–18] Furthermore, in the United States, increases in incidence among blacks of both sexes have outpaced increases among

whites (Figure 1, top graph).^[19] Most recent data (1992–1999) from the National Cancer Institute's Surveillance, Epidemiology and End Results (SEER) Program suggest a leveling off or decrease in this trend for most racial/ethnic groups, with the exception of American Indians in New Mexico (Figure 1, bottom two graphs).^[19,20] Other analyses in the United States have suggested a continued increase among Hispanic women in recent years.^[21] Small numbers of cases during this short time period may or may not indicate a broader trend.

Globally, there is an eightfold variation in kidney cancer incidence.^[17] Racial/ethnic differences within the United States are also significant. According to SEER data, the highest incidence occurs among American Indians in New Mexico (12.3 per 100,000), and the lowest incidence occurs among Asian/Pacific Islanders in California (2.9 per 100,000)

TABLE 1
Incident cases and age-adjusted rates for renal cell carcinoma by race/ethnic group and sex,
NCI-SEER program and California, 1995–1999

	Total			Male			Female		
	Cases	AAIR	95% CI	Cases	AAIR	95% CI	Cases	AAIR	95% CI
Whites (11 SEER Registries) ^a	9071	6.6	6.5–6.8	5719	9.4	9.1–9.6	3352	4.4	4.3–4.6
Non-Hispanic Whites (California) ^c	6085	6.3	6.1–6.4	3922	9.0	8.7–9.3	2163	4.1	3.9–4.3
Blacks (11 SEER Registries) ^a	1181	8.2	7.8–8.7	713	11.5	10.7–12.5	468	5.8	5.2–6.3
Non-Hispanic Blacks (California) ^c	632	7.7	7.1–8.4	409	11.6	10.4–12.8	223	4.8	4.2–5.5
Hispanics (11 SEER Registries) ^b	1009	5.9	5.5–6.3	596	7.9	7.2–8.6	413	4.3	3.9–4.8
New Mexico SEER only ^b	241	10.0	8.7–11.4	140	12.2	10.2–14.6	101	8.0	6.5–9.7
Hispanics (California) ^c	1445	6.2	5.9–6.5	868	8.2	7.6–8.8	577	4.6	4.2–5.0
American Indians (11 SEER Registries) ^a	75	6.5	5.1–8.3	52	10.2	7.5–13.7	23	3.6	2.3–5.5
New Mexico SEER only ^a	52	12.3	9.1–16.4	34	18.7	12.8–27.3	18	7.4	4.3–12.1
Asian/Pacific Islanders (11 SEER Registries) ^a	569	3.6	3.3–3.9	370	5.2	4.7–5.8	199	2.3	2.0–2.6
Asian/Pacific Islanders (California) ^c	407	2.9	2.7–3.3	274	4.4	3.9–5.0	133	1.8	1.5–2.1

AAIR, average annual age-adjusted incidence rate per 100,000 using the 2000 U.S. Standard with 95% confidence intervals (95% CI).

^aData derived from SEER 11 Registries Public Use File November 2001 Submission for Expanded Races 1992–1999 includes 3 of 10 California regional registries.

^bData derived from SEER 11 Registries Public Use File November 2001 Submission for Hispanics 1992–1999 includes 3 of 10 California regional registries.

^cData derived from California Cancer Registry for cases reported as of August 2002.

(Table 1). Relative to the general U.S. population, higher rates have also been noted among American Indian and Alaska Native populations in other geographic areas.^[22–25] Compared with non-Hispanic whites, Hispanics in New Mexico and black populations also have elevated renal cell cancer rates (Table 1). Racial misclassification, as well as differences in methodology used by cancer registries to determine race/ethnicity, account for some portion of the differences by registry for the same racial/ethnic group shown in Table 1.^[22,23]

The underlying reasons for both the upward trends and the racial/ethnic disparity in RCC incidence remain in doubt. Although part of this increase may be attributed to inadvertent clinical detection due to the increased use of ultrasonography, computed tomography, and magnetic resonance imaging, the increasing incidence of distant and unstaged cases of RCC suggest that this increase is not solely due to earlier detection.^[16]

KNOWN RISK FACTORS

Several case-control studies have been conducted to date that examined associations between lifestyle and environmental factors and renal cancer (Table 2). These studies identified a number of risk factors, most notably cigarette smoking, obesity, and hypertension. Smoking and body mass index

(BMI) have demonstrated the most consistent associations with kidney cancer. The strongest association between an exposure and cancers of both the renal parenchyma and pelvis has been with cigarette smoking.^[4,5,26–28] Most case-control and cohort studies have reported significant associations with cigarette smoking, with relative risks ranging from 30% to twofold, and significant dose-response trends.^[5,27] The absence of significant association in a few studies may be partly explained by small sample size and the use of hospital controls with a high prevalence of smoking.^[29–34] Population-based attributable risks for cigarette smoking have ranged from 27% to 37% of RCCs in men and from 10% to 24% in women.^[5,35,36] Consistent positive associations between RCC and obesity have been reported among both men and women.^[6,7,37] The risks tend to rise with increasing BMI. Risks for obese persons have ranged from 20% to nearly threefold, with risks generally higher for women than men.^[37] Assuming a causal association, attributable risks estimated from a population-based case-control study conducted in Minnesota determined that approximately 20% of RCC cases are attributable to excess weight.^[36] Obesity may contribute to risk by increasing levels of endogenous estrogens^[38–40] and availability of insulin-like growth factors.^[41,42] Obesity is also related to hypertension and arterionephrosclerosis, factors that

TABLE 2
Population-based case-control studies of renal cell cancer

Setting/dates of diagnosis	Study/studies	Cases/controls [‡]	Exposures analyzed or controlled for
Australia: New South Wales 1977–1982	[28]	360/985	S, UC, AG, AHY, diet, C, milk, EDU, SES
Australia: New South Wales 1989–1990	[5,10,53]	489/523	S, BMI, HYPT, UC, MC, AHY, DU, DP, OC, HRT, MAR, EDU, OCCU, SROC
Canada: Montreal 1979–1985	[81]	142/533	S, BMI, INC, OCCU, JEM
Canada: 8 provinces 1994–1997	[86]	1279/5370	S, BMI, A, diet, INC, EDU, SES, OCCU, SROC
Canada: Ontario 1986–1987	[4]	518/1381	S, BMI, HBP, DIAB, UC, DU, AG, FH, OC, RH, HRT, C, T, A, diet, ETHN
China: Shanghai 1987–1989	[69]	154/157	S, BMI, HYPT, DU, AG, OC, HRT, RH, diet, EDU, OCCU
Denmark 1989–1992	[6,11,57,222]	365/396	S, TOB, BMI, FH, C, T, A, diet, EDU, SES
Finland 1977–1978	[87]	338/338	S, BMI, C, OCCU, JEM
Germany: West Berlin, Bremen, Leverkusen, Halle, and Jena 1991–1995	[88]	935/4298	S, BMI, SES, OCCU, JEM
Germany: Rhein-Neckar-Edenwald 1989–1991	[94]	277/286	S, SES, MAR, urban, OCCU, SROC
International Renal Cell Cancer Study [‡] 1989–1991	**	1732/2309	S, TOB, BMI, HT, WT, PA, HYPT, DIAB, UC, FH, MC, AHY, DU, AG, DP, RH, HYST, OC, HRT, A, diet, EDU, OCCU, SROC
Sweden 1996–1998	[223]	648/900	S, BMI, HYPT, DIAB, HT, BWT, EDU
Sweden 1989–1991	[224]	379/353	S, BMI, HT, WT, WT change, DP, PA
USA: Minneapolis-St. Paul, Minnesota 1974–1979	[35,90]	495/691	S, BMI, UC, AG, C, T, A, diet, ETHN, OCCU
USA: Minnesota 1988–1990	[64,117]	438/687	S, BMI, HYPT, DU, AHY, OCCU, JEM
USA: Boston Metro Area 1981–1984	[77]	518/518	S, BMI, drugs, VIT, PA, MC, diet, ETHN, OCCU
USA: Boston Metro Area 1976–1983	[68]	203/605	S, BMI, HYPT, AG, UC, MC, diet, ETHN, EDU, INC
USA: Iowa 1986–1989	[225–227]	406/2429	S, BMI, HYPT, BI, FH, PA, C, T, A, diet, OCCU
USA: Los Angeles County/1975–1979	[7]	160/160	S, BMI, HYPT, UC, DP, DU, C, T, A, HRT, RH, MAR, REL, EDU, OCCU, SROC
USA: Los Angeles County 1986–1994	[228–231]	1204/1204	S, BMI, HYPT, FH, AG, DIAB, MC, DP, DU, AHY, HYST, A, diet, EDU, OCCU
USA: Western Washington State 1996–1997	[217]	130/505	S, BMI, HBP, GST

A, alcohol; AG, analgesics; AHY, antihypertensive drugs; BMI, body mass index; BWT, birth weight; C, coffee; DIAB, diabetes; DP, diet pills; DU, diuretics; EDU, education; ETHN, ethnicity; FH, family history of kidney cancer; GST, glutathione S-transferase polymorphisms; HBP, high blood pressure; HRT, hormone replacement therapy; HT, height; HYPT, hypertension; HYST, hysterectomy; INC, income; JEM, job exposure matrix; MAR, marital status; MC, medical conditions; OCCU, occupation; PA, physical activity; REL, religion; RH, reproductive history; SES, socioeconomic status; SROC, self-reported occupational compounds; T, tea; TOB, tobacco; UC, urologic conditions; VIT, vitamins; WT, weight.

[‡]Number of cases and controls may differ by analyses.

**=Refs. [27,49,70,79,94,232–234] 16 centers in 5 countries: Australia (Sydney), Denmark (Denmark), Germany (Berlin and Heidelberg), Sweden (Uppsala), and USA (Minnesota).

could alter the renal cell environment, making it favorable to tumor induction.^[43,44] Although obesity is a stronger and more consistent risk factor in women than men, it has been suggested that female sex hormones may reduce RCC risk. A history of oophorectomy/hysterectomy is associated with an increased risk of renal cell cancer.^[45–47] Similarly, testosterone treatment and/or oophorectomy increase the incidence of RCC, whereas estradiol or estriol treatment and/or castration decrease the incidence of RCC in rats treated with the renal carcinogen ferric nitrilotriacetate (Fe-NTA).^[48]

Hypertension has been associated with RCC in most studies, with excess risks ranging from 20% to threefold.^[49–52] It remains unclear whether hypertension or treatment of hypertension is the main risk factor. Some studies have found that risks remain elevated after adjustment for antihypertensive medications,^[50,53–56] others have found that risks dropped to insignificant levels,^[4,7,49,57] and others have not adjusted for medications because their use is highly correlated with the disease.^[51,52,58,59] In the United States, national surveys indicate that the prevalence of hypertension in the population has remained relatively stable; however, the number and types of medications used to treat hypertension have increased substantially.^[60] Such contrasting trends between the disease and its treatment stress the importance of investigating the role of various families of antihypertensive medications (diuretics, beta blockers, calcium channel blockers) and RCC incidence because the prevalence of hypertension has not been increasing. Moreover, the long-term effect of consuming hypertensive medications and the vascular and cellular effect on the kidney is unknown.^[50,53,56]

Diabetes, also considered an emerging risk factor for RCC, may be related in part to both hypertension and obesity because obesity contributes to both conditions.^[61] Results from several case-control studies have found little evidence of an association. However, before the study of diabetes as a possible risk factor can be dismissed, assessment of potential confounding by associated conditions will be required before a causal link between diabetes and RCC can be accepted.^[36,61]

As stated earlier, increases in kidney cancer incidence have been reported in the United States and globally.^[16,17] Attempts to explain both the increase in incidence and the racial variation by examining national and global trends of known risk factors have not been successful;^[16] however, the upward trends in the prevalence of these two risk factors have varied little among race and ethnic groups.^[62] Both obesity and hypertension are known risk factors for RCC and are more prevalent among blacks than whites in the United States. Recent evidence from the National Health and Nutrition Examination Survey (NHANES) suggests that age-specific increases in hypertension among black males may be playing a role.^[63]

In the United States, hypertension and obesity may explain in part why RCC incidence is higher among the black population but does not account for the increases in incidence

observed worldwide.^[17] Smoking, an established risk factor for RCC has declined in past decades; however, the long latency of RCC could explain some but not all of the cases in older age groups. Lastly, in the United States, the largest increases in RCC have occurred for early-stage tumors. A proportional increase in early-stage tumors compared with advanced tumors could be explained by improvements in detection. However, better detection does not explain the increases observed in advanced tumors and RCC mortality.^[16] This trend of rising incidence and mortality has been observed among U.S. blacks compared to U.S. whites and in several other developed countries, suggesting that risk factors are contributing to this disease.

Other risk factors that are associated with renal cancer include abuse of analgesics^[4,8,57,64] and history of cystic disease due to long-term hemodialysis.^[9,65,66] Excess risks have also been associated with kidney stones, cysts, and infection; however, the findings have not been consistent.^[7,27,28,35,67] The role of diet has been investigated, and excess risks have been observed with consumption of certain meats and a reduced risk with certain fruits and vegetables.^[29,32,68–70] Wolk et al. observed significant positive associations with total energy intake (RR 1.7; 95% CI 1.4–2.2) in the highest vs. the lowest quintile of exposure.^[70] Fried meats were also associated with increased RCC risk, whereas vegetables and fruits were protective. Increased risk was also observed in those with the lowest intake (lowest decile) of vitamin E and magnesium intake. Lindbland et al. in 1997 also observed increased risk with frequent intake of fried or sautéed meat or poultry consumption, and a significantly protective effect on RCC risk was observed with increased consumption of fruit, especially citrus fruits.^[71] Yuan et al. observed strong inverse trends between cruciferous and dark green vegetable intakes and RCC risk (p-trend<0.001).^[72]

OCCUPATIONAL EXPOSURES

Although occupational cohort studies comprise the bulk of epidemiological literature examining employment-related kidney cancer risk, several population-based RCC case-control studies have been conducted and generally adjust occupational risk estimates for the effects of confounders such as smoking and often BMI (Table 2). Although RCC has not generally been considered an occupational cancer, several occupationally derived exposures have been implicated in a growing body of research, including asbestos, gasoline fumes, chlorinated solvents, diesel exhaust, polycyclic aromatic hydrocarbons (PAHs), printing and dyes, cadmium, and lead.

Asbestos

Asbestos fibers have been shown to induce kidney cancers in animals, and asbestos bodies can occur in kidneys of persons with asbestosis.^[73–75] Several epidemiological studies have reported elevated risks among asbestos workers, shipyard

workers, insulation workers, and seafarers, as well as persons occupationally exposed to asbestos.^[10,11,73,76–83] Some population-based case-control studies have found significantly elevated risks ranging from 1.4 to 1.6.^[10,79] However, other population-based case-control studies and two meta-analyses of occupational cohorts with asbestos exposure have not found increased risks.^[35,84–87] Positive trends with intensity^[88] or duration^[79–83] of exposure have not been observed. McCredie et al.^[10] observed higher risks associated with the year asbestos exposure began (OR=5.28, 95% CI: 1.43–19.55; 1956–1986 vs. 1929–1948). Although elevated, exposure in this study was self-reported and may have been subject to recall bias.

Automotive Gasoline Fumes

Following a study demonstrating renal cancer among rats chronically exposed to unleaded gasoline fumes,^[35,89,90] McLaughlin identified elevations in RCC risk associated with increased duration of employment among gas station attendants.^[90] In addition, cohort and case-control studies among this group of workers have reported a 20–40% elevation in risk ratios, although not all findings were statistically significant.^[79,90–92] Occupational gasoline exposure, both self-reported^[79] and JEM-estimated,^[87] has been significantly associated with RCC in two case-control studies. Other case-control studies have found no association for gasoline^[7,58,93] or petroleum product exposure.^[94] Partanen et al.^[87] identified a dose-response relationship with respect to the estimated (ppm) and cumulative (ppm-years) exposure level (OR=4.3; 95% CI: 1.2–16.4; 14–102 ppm-years of exposure compared to controls without occupational exposure). However, an association was not observed with duration of job exposure alone. Hu et al. found a dose-response increase for self-reported (occupational and other) benzene exposure, attributing the majority of occupational benzene exposure to petroleum products.^[86] Other studies found no elevation in risk.^[7,81,95,96] Conflicting findings may be due to low statistical power to detect an association in the smaller case-control studies, difficulty in assessing the intensity of exposure from job title alone, geographic variation in gasoline constituents, and decrease in daily dose to gasoline attendants over time with the phase out of full-serve pumps.^[97]

Solvents

Elevated rates of kidney cancer have been observed among dry cleaners, aerospace workers, aircraft maintenance workers, and architects^[10,55,98–104] although not in all studies.^[105–107] Schlehofer et al.^[94] observed an increased risk of RCC among men reporting exposure to chlorinated solvents (OR=2.5; 95% CI: 1.2–5.2). Mandel et al.^[79] reported an increase in RCC among men who reported they were ever occupationally exposed to dry cleaning solvents (OR=1.4; 95% CI: 1.1–1.7). However, when duration of occupation was considered, men in the midrange of exposure (8–25 years of exposure) had the highest level of risk.

TCE has been of particular interest due to widespread industrial use as a degreasing agent and increasing awareness of nonoccupational exposure pathways.^[108] Most TCE-exposed occupational cohorts have not demonstrated statistically significant increases in risk.^[103,104,106,109–112] Although not statistically significant, aerospace workers with airborne TCE exposures above 50 ppm were at a near twofold risk of kidney cancer mortality (RR=1.9; 95% CI 0.9–4.2, adjusted for age and sex), compared with workers exposed to lower levels.^[103] Additional exposures to workers drinking and showering in contaminated plant well water (730–2200 ppb of TCE) were not considered.^[103] Two studies have found very high risks for occupational TCE exposure,^[113,114] although their validity has been questioned.^[106,115] A subsequent hospital-based case-control study by the same research group found a significant association with self-reported TCE exposure and consequent narcotic symptoms.^[116]

Data on TCE from population-based case-control studies are limited. Using a JEM, Pesch et al.^[88] found a nonsignificant elevated risks for TCE, perchloroethylene (PCE), and all chlorinated solvents. Dosemeci et al.^[117] found significantly higher (twofold) risks for renal cell cancer among women exposed to chlorinated solvents, as well as TCE, suggesting that women may be at increased risk.^[117,118] Several other studies have found higher risk ratios among women compared with men, although they were not specifically designed to examine gender differences.^[11,55,79,117,119,120]

Biological plausibility for the TCE/RCC link has also been derived from animal studies of TCE exposure and RCC induction, epithelial damage to the proximal renal tubules following high levels of exposure to chlorinated solvents in humans, and the more recent finding of a C to T nucleotide transition on codon 81 of the VHL gene.^[121] Bruning^[122] has suggested a threshold determination for TCE metabolism in humans. At low-dose exposure, the oxidative metabolic pathway is activated. At higher exposures, this pathway becomes saturated, and TCE is metabolized via glutathione conjugation, leading to the formation of damaging reactive metabolites in the proximal tubule.^[122,123]

Diesel Exhaust

Occupational cohort and case-control studies have found elevated risks for RCC among truck drivers and urban bus drivers.^[11,55,67,116,124,125] Brownson^[67] found a significant threefold risk for heavy truck drivers (OR=3.1; 95% CI: 1.1–8.5). Using a job exposure matrix, Boffeta et al.^[124] found an elevated risk for exposure to diesel exhaust in a Swedish occupational cohort, although this study did not adjust for possible confounding effects of smoking. In a population-based case-control study, Schlehofer et al. found a significantly elevated risk among men reporting occupational exhaust exposure for at least 5 years (OR=1.8; 95% CI: 1.0–3.2).^[94] Constituents of diesel and motor exhausts of particular interest include polycyclic aromatic hydrocarbons (PAHs).

Polycyclic Aromatic Hydrocarbons

Elevated RCC risks among coke oven workers and petroleum refinery workers in occupational cohort studies have generated interest in PAH exposure.^[79,126] Among coke oven workers, a highly increased risk, relative to other steel workers, was not replicated in the same cohort after a 30-year follow-up study.^[126–128] Elevated risks in occupational cohort studies and nested case-control studies of petroleum refinery workers have not been consistent.^[128–137] Other studies of occupational cohorts highly exposed to PAHs (including aluminum reduction) have found elevated but not statistically significant kidney cancer risks.^[138–141] County-level kidney cancer mortality and the proportion of the population employed in petroleum refining and related industries have historically shown a crude ecologic correlation in the United States.^[142] Population- and hospital-based case-control studies have found elevated risks for employment in the oil refinery industry,^[55,79,90] and self-reported exposure to burning coal, petroleum, tar and/or pitch products,^[35,86,95] with two of these studies providing a suggestive dose-response effect with duration of employment^[35] and intensity of exposure.^[95] Two population-based case-control studies that used a job exposure matrix to estimate PAH intensity found no association or dose-response effect.^[87,88]

Printing and Dyes

Elevated RCC risks have been identified for printers, newspaper pressmen, paperboard printing workers, color film developers, as well as for individuals exposed to paints, pigments, and inks.^[81,88,143–145] Other studies have found no association with printing industry employment.^[10,11] Occupational cohorts exposed to dyes containing lead and cadmium demonstrated elevated kidney cancer mortality.^[146,147] JEM-based exposure to chromium (VI) compounds, also used in inorganic dyes, leather tanning, and wood preserving, was associated with increased risk in one case-control study,^[81] although concurrent exposure to lead and cadmium was not accounted for. Tanners and textile workers have shown elevated risks.^[148,149] Self-reported occupational exposure to benzidine was associated with increased risk, although studies among cohorts exposed to benzidine have not clearly demonstrated an association with kidney cancer.^[86]

Cadmium

Three major sources of cadmium exposure include diet, cigarette smoking, and occupation. Kolonel reported a crude association with occupational cadmium exposure.^[150] Population-based case-control studies by Mandel et al.^[79] and Hu et al.^[86] observed a significantly elevated risk for self-reported exposure to cadmium and cadmium salts among men. Using job exposure matrices, Pesch et al.^[88] observed excess risks for high exposure (OR=1.4; 95% CI: 1.1–1.8 men; OR=2.5; 95% CI: 1.2–5.3 women), and Partanen et al.^[87] found elevated risk with any exposure (OR=4.4; 95% CI: 0.4–43.0),

although based on only three exposed cases. Other case-control and occupational cohort studies have found no evidence of association.^[10,35,151–154] Evidence of a dose-response pattern with increasing exposure intensity or duration of employment has not been found.^[79,88]

Lead

Lead has been shown to induce renal cancers in animals and nephropathy among humans with high occupational exposures.^[155–157] Three occupational cohort studies of lead smelter work have found a 1.4- to threefold elevation in kidney cancer mortality, depending on job area/task,^[158–160] although only one demonstrated statistically significant findings.^[158] Case-control studies have found significantly elevated risks among welders.^[10,81,88,161–163] Welding fumes have been associated with an increased frequency of VHL gene mutations in RCC patients in a case-case comparison.^[164] JEM-assessed lead exposure has been significantly associated with RCC in two case-control studies.^[87,88] Studies of other lead exposed cohorts and meta-analyses of lead exposed occupational cohorts have found no statistically significant excesses of kidney cancer.^[157,165,166]

Other Occupational Associations

Other industries and occupations associated with elevations in RCC risk include iron and steel industries,^[87,88,163,167] metal foundries,^[168] electrical utilities,^[169,170] electronic equipment assembly,^[88] electricians and electrical maintenance,^[171] aircraft mechanics,^[81,104] and firefighters.^[172,173] There have been other associations reported among airline pilots,^[174] farmers,^[81] physicians,^[166] pulp and paper mill workers,^[175] railway workers,^[176] miners exposed to dinitrotoluene,^[177] rubber workers,^[88] copper smelter work,^[178] plumbers and pipefitters,^[179] and metal furniture factory work.^[180] Other occupational exposures investigated by a few studies include chromium (VI),^[81,116] ozone,^[81] jet/aviation fuel,^[81,93] UV radiation,^[81] styrene-butadiene rubber,^[81] heat stress,^[139] and copper sulfate.^[181]

ENVIRONMENTAL EXPOSURES

There are currently few investigations of renal cancer and carcinogenic or infectious disease agents in air, water, or food. Research to date has focused on exposures to chlorination by-products, arsenic and asbestos, in household drinking water supplies.

In an ecological study, Cantor et al. reported an increasing correlation with residual mortality (i.e., the proportion of mortality unexplained by industry, age, urban residence) as the level of trihalomethanes in drinking water of counties increased.^[182] Wilkins and Comstock found no evidence of an association.^[183] Yang et al. matched chlorinating and nonchlorinating communities on level of rurality and found a significantly increased risk among chlorinating communities

for both males and females.^[184] In a population-based case-control study, Koivusalo et al. used historical (1950–1987) measures of turbidity and biotic components of public drinking water in an equation to estimate the level of mutagenicity in drinking water.^[185] Men exposed to 3000 net revertants/litre were more likely to have developed renal cancer (OR=1.49; 95% CI: 1.05–2.73), although a significant increase was not observed in women. Missing data for individual smoking history (43%) limited the ability of this study to adjust for possible confounding.

Several ecological studies of arsenic exposure have found elevated rates of kidney cancer in areas with high exposure to arsenic in groundwater sources.^[181–190] Dose-response patterns are evident in the studies by Wu et al.^[187] and Hopenhayn-Rich.^[189] In a case-control study of a rural agricultural population, Kurtzio et al. measured arsenic levels in drilled (not dug) wells and did not find an elevated risk of kidney cancer, although this study was limited by a small number of cases and relatively low-exposure levels.^[191]

Exposure to asbestos in drinking water has been of interest due to the use of asbestos-cement pipe for transporting public water and exposure in public surface water supplies due to runoff from mine tailings. Two ecological studies of household drinking water have found significantly elevated risks for kidney cancer in men but not women^[192,193] and another found only weak associations among women.^[194] One study reported elevated kidney cancer mortality among women residing in counties containing an asbestos mine.^[195] These studies lack information on residential history and water use patterns. A major issue has been the assumption of past exposure based on current measurements.^[196] Moreover, dose-response patterns have not been investigated.

GENETIC SUSCEPTIBILITY

Numerous reports of familial RCC have thus far identified four genes that are associated with several inherited forms of kidney cancer.^[197] Tumors associated with the Von Hippel Lindau (VHL) syndrome are primarily of the clear cell type and are associated with germline mutations of the *VHL* gene located on chromosome 3p.^[198] VHL is a tumor suppressor gene (3p25–p26) whose protein product forms a transcription-regulatory complex targeting the α -subunits of hypoxia-inducible factor (HIF) for ubiquitin-mediated degradation in an oxygen-dependent manner.^[199] Loss of VHL protein (pVHL) activity can fully activate HIF-mediated responses by stimulating the expression of hypoxia-sensitive genes involved in energy metabolism, angiogenesis, hematopoiesis, and oxygen delivery. Mutations within the β -subunit of pVHL disrupt its interaction with HIF- α subunits leading to constitutive HIF- α stabilization. Striking up-regulation of HIF target genes in tumors that lack functional pVHL demonstrates the importance of this protein in the regulation of the HIF transcriptional cascade mediated through cellular

responses to hypoxia.^[199–203] Familial occurrence of RCC also has been described in the absence of the inherited VHL syndrome and has featured constitutional (3:8t) balanced translocations involving chromosome 3p.^[204] In familial cases of VHL disease, patients inherit a defective tumor suppressor gene in every cell but do not develop tumors until the normal allele becomes mutated or deleted (somatic loss). In sporadic RCC, somatic VHL mutations have been observed in about 50% of tumors.^[205,206] Epigenetic silencing of the gene through promoter methylation has been observed in approximately 20% of sporadic tumors.^[207,208] These observations suggest that the VHL gene may be inactivated in as many as 70% of sporadic RCC tumors.

Because the VHL gene is frequently mutated in RCC, mutational spectrum analysis has been used to look for carcinogen-specific mutational fingerprints (Table 3). Bruning et al.^[209] and Brauch et al.^[121] provided the first possible link between trichloroethylene (TCE) and specific somatic mutations in the VHL gene in tumor DNA from highly exposed workers. A high frequency of VHL gene mutation was reported among patients with high cumulative exposure to TCE, but not among unexposed patients. More recent investigations have proposed additional hotspot mutations in this gene.^[206,210] Specific types of VHL mutation have also been associated with the GSTT1 active and NAT1 slow/rapid genotypes.^[205] Although the VHL mutational studies conducted to date provide promising leads for the existence of mutational hotspots among potentially susceptible populations, a larger study of the VHL mutational spectrum using improved exposure assessment methods and including information on other risk factors and confounding variables will be required to elucidate these findings.

Unlike clear cell renal carcinomas, hereditary papillary renal cancer is related to germline mutations and activation of the MET proto-oncogene located on chromosome 7p.^[211] Unlike the VHL gene, MET oncogene provides the cell with positive growth signals that can lead a cell into unregulated cell cycling, morphogenesis, and eventual tumorigenesis. MET gene mutations give rise to kidney cells with mutant hepatocyte growth factor (HGF) receptors. Unlike cells with normal receptors, these cells are unable to turn off from the activated state after HGF binding. HGF is expressed in many tissues and appears to be a cellular mitogen.^[212] It is noteworthy that somatic mutations in both MET and VHL genes have been described in a high proportion of sporadic cases of papillary and clear cell cancers, respectively, implicating that alterations in these genes could play a major role in RCC development in the general population. Aside from TCE-associated VHL mutations, it is currently unknown whether specific lifestyle factors or exposures alter these genes directly.

Recent reports have identified additional genes involved in two rare inherited forms of renal cancer. The fumarate hydratase (FH) gene located on chromosome 1q42-3-q43 was recently discovered to be associated with a certain form of

TABLE 3
Mutational spectrum studies of the von Hippel-Lindau (VHL) gene and renal cell cancer (RCC)

RCC study size	Types of VHL mutations observed	Reference	Exposures
173	42% of cases mutated 4 polymorphisms identified @ nucleotide(nt) 194 (codon 65), nt 27 (codon 9), nt 183 (codon 61), nt 435 (codon 145)	[220]	Not known
227	45% of cases mutated 12% ATT.TTT at codon 147/148 Mutation prevalence was associated with stage	[210]	Not known
102	54% of cases mutated in clear cell RCC 18% of cases mutated in chromophilic tumors 6% in codon 160 5% in codon 41–45	[206]	Not known
102 (same study group as above)	Welding fumes associated with increased number of multiple mutations (OR=5.65; 95% CI: 1.39–22.93) Citrus-associated with fewer G to A transitions (OR=0.13; 95% CI: 0.03–0.57) Selenium associated with fewer G to A transitions (OR=0.31; 95% CI: 0.10–1.02) and multiple mutations (OR=0.27; 95% CI 0.07–0.97) Smoking was not a risk factor for mutations but appeared to change the outcome of many variables	[164]	Occupational exposures, lifestyle and nutritional factors while controlling for possible confounding factors
195	46% of cases mutated No hotspot observed but more tranversions occurred in cases carrying a GSTT1 active allele and in those with NAT slow/rapid genotypes	[205]	Not known
23	100% of cases mutated 30% in exon 1 44% in exon 2 26% in exon 3	[209]	Trichloroethylene
44	75% of cases mutated 52% in exon1 20% in exon 2 28% in exon 3 39% contained a hotspot mutation at nt 454 (codon 81)	[121]	Trichloroethylene

familial papillary kidney cancer.^[213,214] Individuals with aberrant FH genes are also predisposed to several other dominantly inherited conditions such as uterine fibroids and skin leiomyomata. Birt-Hogg-Dubé syndrome (BHD) is another inherited form of kidney cancer that has been under study. This syndrome is primarily associated with chromophobe RCCs although clear cell tumors and hybrid chromophobe/oncocytic tumors are also commonly found.^[197] The gene associated with BHD syndrome, located on chromosome 17p, is dominantly inherited.^[215] Patients af-

flicted with BHD are also susceptible to cutaneous fibrofolliculomas and trichodiscomas on the face, neck, back, and chest; lung cysts, spontaneous pneumothorax; and multifocal or bilateral renal cancers. It is unknown whether alterations in either of these genes are observed in sporadic tumors.

Also in the genetic arena, several studies have demonstrated that individuals carrying polymorphisms in certain phase I and II metabolism genes may be at increased risk of RCC. Schultz et al. observed a 70% elevated risk of RCC in

individuals carrying the NQO1 homozygous null-allele vs. wild-type genotype.^[216] Sweeney et al.^[217] observed an excess of individuals carrying the inactive GSTT1 null polymorphism in cases compared with controls (OR=1.9; 95% CI: 1.1–3.4). This relationship was modified by BMI, and among those in the lowest tertile, the OR for RCC increased to 4.8 (95% CI: 1.8–13.0). Bruning et al. reported that the odds for TCE exposure in cases compared with controls were 2.7 (95% CI: 1.18–6.33) for individuals carrying the active GSTM1 polymorphism and 4.2 (95% CI: 1.16–14.91) for those carrying the active GSTT1 polymorphism.^[218] Semenza et al.^[219] reported that subjects carrying the slow acetylator polymorphisms of N-acetyltransferase 2 gene (NAT2) had a twofold increased risk of RCC (OR=1.8; 95% CI: 1.1–2.9). Although the OR for RCC in smokers was 2.2 (95% CI: 1.3–3.7), the risk was higher among smokers carrying the slow acetylator polymorphism (OR=3.2; 95% CI: 1.7–6.1) than among smokers carrying the rapid acetylator polymorphism (OR=1.4; 95% CI: 0.7–2.9). Longuemaux et al.^[220] found the CYP1A1 variant allele to be associated with a 2.1-fold (95% CI: 1.1–3.9) increase in RCC risk. A higher risk was also observed in subjects carrying the CYP1A1 variant polymorphism when combined with the active GSTT1 polymorphism (OR=2.3; 95% CI: 1.2–4.5), the NAT2 slow acetylator polymorphism (OR=2.6; 95% CI: 1.2–5.8), and the CYP1A1 (OR=3.5; 95% CI: 1.1–11.2). Although these studies show promising hypotheses that need to be further explored, most of the studies conducted to date have been too small for analyses of gene-environment interactions (study size ranged from 50 to 220 cases). Most of these studies also lacked important information regarding other risk factors and potential confounding factors.

CONCLUSION

Results from currently available epidemiological studies provide convincing evidence linking smoking, obesity, and hypertension to an elevated risk of RCC, potentially accounting for 50% of all U.S. cases. The variation in the prevalence of these factors plus other related comorbidities, such as diabetes, across subpopulations may help to explain at least a portion of the racial and geographic variation in RCC incidence observed, not only in the United States but worldwide. The multifactorial nature of RCC requires that additional work be conducted to explain risks associated with individual factors and to elucidate complex relationships between potential genetic, lifestyle, and environmental elements on cancer development. Recently, a new unifying mechanism for the etiology of RCC has been described which unites the many risk factors associated with RCC through among different lines of research. Lipid peroxidation has been suggested as a unifying mechanistic pathway by which several risk factors, including obesity, hypertension, and chemical exposure-induced lipid peroxidation of the proximal renal

tubules, could induce renal carcinogenesis.^[221] This hypothesis may also explain the roles of other risk factors related to hormone status, smoking, diabetes, and protective factors such as dietary antioxidants for RCC. Although an interesting hypothesis, this possible causal mechanism has not yet been tested in vivo.

Because approximately 50% of sporadic RCC cases remain unexplained by established risk factors, more relationships need to be elucidated with respect to potential occupational and environmental exposures and cancer development. This review article covered those risk factors for which the strongest associations in humans have been noted. Epidemiological studies often have a limited ability to establish causality due to an inconsistent case definition, imprecise measures of exposure (years employed, job title, or unspecified chemical exposures), and a lack of consistent control for confounding factors (smoking, comorbidities, or other lifestyle factors). The application of biological markers of genetic susceptibility and exposure as well as intermediate biological endpoints that target RCC and conditions associated with increased risk could help to identify related factors and reduce the observed disparity in subpopulations at greatest risk. Future studies need to continue addressing these study design issues, in addition to considering underlying factors that may influence individual risk and help explain gender and racial variation of RCC.

REFERENCES

1. Landis, S.H.; Murray, T.; Bolden, S.; Wingo, P.A. Cancer statistics, 1999. *CA Cancer J. Clin.* **1999**, *49* (1), pp. 1, 8–31.
2. American Cancer Society. *Cancer Facts and Figures, 2003*; American Cancer Society: Atlanta, GA, 2003.
3. Lynch, C.F.; Cohen, M.B. Urinary system. *Cancer* **1995**, *75* (Suppl. 1), 316–329.
4. Kreiger, N.; Marrett, L.D.; Dodds, L.; Hilditch, S.; Darlington, G.A. Risk factors for renal cell carcinoma: results of a population-based case-control study. *Cancer Causes Control* **1993**, *4* (2), 101–110.
5. McCredie, M.; Stewart, J.H. Risk factors for kidney cancer in New South Wales. I. Cigarette smoking. *Eur. J. Cancer* **1992**, *28A* (12), 2050–2054.
6. Møller, A.; Engholm, G.; McLaughlin, J.K.; Olsen, J.H. Risk factors for renal-cell carcinoma in Denmark. III. Role of weight, physical activity and reproductive factors. *Int. J. Cancer* **1994**, *56* (1), 66–71.
7. Yu, M.C.; Mack, T.M.; Hanisch, R.; Cicioni, C.; Henderson, B.E. Cigarette smoking, obesity, diuretic use, and coffee consumption as risk factors for renal cell carcinoma. *J. Natl. Cancer Inst.* **1986**, *77* (2), 351–356.
8. McCredie, M.; Pommer, W.; McLaughlin, J.K.; Stewart, J.H.; Lindblad, P.; Mandel, J.S.; Møller, A.; Schlehofer, B.; Niwa, S. International renal-cell cancer study. II. Analgesics. *Int. J. Cancer* **1995**, *60* (3), 345–349.
9. Bretan, P.N., Jr.; Busch, M.P.; Hricak, H.; Williams, R.D. Chronic renal failure: a significant risk factor in the development of acquired renal cysts and renal cell carcinoma. Case reports and review of the literature. *Cancer* **1986**, *57* (9), 1871–1879.
10. McCredie, M.; Stewart, J.H. Risk factors for kidney cancer in New

- South Wales. IV. Occupation. *Br. J. Ind. Med.* **1993**, 50 (4), 349–354.
11. Møller, A.; Engholm, G.; McLaughlin, J.K.; Olsen, J.H. Occupational risk factors for renal-cell carcinoma in Denmark. *Scand. J. Work, Environ. & Health* **1994**, 20 (3), 160–165.
12. Linehan, W.M.; Lerman, M.I.; Zbar, B. Identification of the von Hippel-Lindau (VHL) gene. Its role in renal cancer. *JAMA* **1995**, 273 (7), 564–570.
13. Schmidt, L.; Duh, F.M.; Chen, F.; Kishida, T.; Glenn, G.; Choyke, P.; Scherer, S.W.; Zhuang, Z.; Lubensky, I.; Dean, M.; Allikmets, R.; Chidambaram, A.; Bergerheim, U.R.; Feltis, J.T.; Casadevall, C.; Zamarron, A.; Bernues, M.; Richard, S.; Lips, C.J.; Walther, M.M.; Tsui, L.C.; Geil, L.; Orcutt, M.L.; Stackhouse, T.; Zbar, B. Germline and somatic mutations in the tyrosine kinase domain of the MET proto-oncogene in papillary renal carcinomas. *Nat. Genet.* **1997**, 16 (1), 68–73.
14. Weirich, G.; Glenn, G.; Junker, K.; Merino, M.; Storkel, S.; Lubensky, I.; Choyke, P.; Pack, S.; Amin, M.; Walther, M.M.; Linehan, W.M.; Zbar, B. Familial renal oncocytoma: clinicopathological study of 5 families. *J. Urol.* **1998**, 160 (2), 335–340.
15. Pavlovich, C.P.; Walther, M.M.; Eyler, R.A.; Hewitt, S.M.; Zbar, B.; Linehan, W.M.; Merino, M.J. Renal tumors in the Birt-Hogg-Dube syndrome. *Am. J. Surg. Pathol.* **2002**, 26 (12), 1542–1552.
16. Chow, W.H.; Devesa, S.S.; Warren, J.L.; Fraumeni, J.F., Jr. Rising incidence of renal cell cancer in the United States. *JAMA* **1999**, 281 (17), 1628–1631.
17. Mathew, A.; Devesa, S.S.; Fraumeni, J.F., Jr.; Chow, W.H. Global increases in kidney cancer incidence, 1973–1992. *Eur. J. Cancer Prev.* **2002**, 11 (2), 171–178.
18. Ries, L.A.G. *SEER Cancer Statistics Review, 1973–1999*, National Cancer Institute; NCI: Bethesda, MD, 2002.
19. SEER 9 Registries. *Public Use File Submission*, NCI, DCCPS, Surveillance Research Program, Cancer Statistics Branch, 2001.
20. SEER 11 Registries. *Public Use File Submission*, NCI, DCCPS, Surveillance Research Program, Cancer Statistics Branch, 2001.
21. Canto, M.T.; Chu, K.C. Annual cancer incidence rates for Hispanics in the United States: surveillance, epidemiology, and end results, 1992–1996. *Cancer* **2000**, 88 (11), 2642–2652.
22. Lanier, A.P.; Alberts, S.R. Kidney and bladder cancer in Inuit 1969–1988. *Acta Oncol.* **1996**, 35 (5), 595–599.
23. Cobb, N.; Paisano, R. Patterns of cancer mortality among Native Americans. *Cancer* **1998**, 83 (11), 2377–2383.
24. Partin, M.R.; Rith-Najarian, S.J.; Slatyer, J.S.; Korn, J.E.; Cobb, N.; Soler, J.T. Improving cancer incidence estimates for American Indians in Minnesota. *Am. J. Public Health* **1999**, 89 (11), 1673–1677.
25. Frost, F.; Taylor, V.; Fries, E. Racial misclassification of Native Americans in a surveillance, epidemiology, and end results cancer registry. *J. Natl. Cancer Inst.* **1992**, 84 (12), 957–962.
26. Wynder, E.L.; Mabuchi, K.; Whitmore, W.F., Jr. Epidemiology of adenocarcinoma of the kidney. *J. Natl. Cancer Inst.* **1974**, 53 (6), 1619–1634.
27. McLaughlin, J.K.; Lindblad, P.; Møller, A.; McCredie, M.; Mandel, J.S.; Schlehofer, B.; Pommer, W.; Adami, H.O. International renal-cell cancer study. I. Tobacco use. *Int. J. Cancer* **1995**, 60 (2), 194–198.
28. McCredie, M.; Ford, J.M.; Stewart, J.H. Risk factors for cancer of the renal parenchyma. *Int. J. Cancer* **1988**, 42 (1), 13–16.
29. Armstrong, B.; Garrod, A.; Doll, R. A retrospective study of renal cancer with special reference to coffee and animal protein consumption. *Br. J. Cancer* **1976**, 33 (2), 127–136.
30. Goodman, M.T.; Morgenstern, H.; Wynder, E.L. A case-control study of factors affecting the development of renal cell cancer. *Am. J. Epidemiol.* **1986**, 124 (6), 926–941.
31. Asal, N.R.; Risser, D.R.; Kadamani, S.; Geyer, J.R.; Lee, E.T.; Cherng, N. Risk factors in renal cell carcinoma. I. Methodology, demographics, tobacco, beverage use, and obesity. *Cancer Detect. Prev.* **1988**, 11 (3–6), 359–377.
32. Talamini, R.; Baron, A.E.; Barra, S.; Bidoli, E.; La Vecchia, C.; Negri, E.; Serraino, D.; Franceschi, S. A case-control study of risk factor for renal cell cancer in northern Italy. *Cancer Causes Control* **1990**, 1 (2), 125–131.
33. Benhamou, S.; Lenfant, M.H.; Ory-Paoletti, C.; Flamant, R. Risk factors for renal-cell carcinoma in a French case-control study. *Int. J. Cancer* **1993**, 55 (1), 32–36.
34. Siemiatycki, J.; Krewski, D.; Franco, E.; Kaiserman, M. Associations between cigarette smoking and each of 21 types of cancer: a multi-site case-control study. *Int. J. Epidemiol.* **1995**, 24 (3), 504–514.
35. McLaughlin, J.K.; Mandel, J.S.; Blot, W.J.; Schuman, L.M.; Mehl, E.S.; Fraumeni, J.F., Jr. A population-based case-control study of renal cell carcinoma. *J. Natl. Cancer Inst.* **1984**, 72 (2), 275–284.
36. Benichou, J.; Chow, W.H.; McLaughlin, J.K.; Mandel, J.S.; Fraumeni, J.F., Jr. Population attributable risk of renal cell cancer in Minnesota. *Am. J. Epidemiol.* **1998**, 148 (5), 424–430.
37. Chow, W.H.; Gridley, G.; Fraumeni, J.F., Jr.; Jarvholm, B. Obesity, hypertension, and the risk of kidney cancer in men. *N. Engl. J. Med.* **2000**, 343 (18), 1305–1311.
38. Kumar, A.; Mittal, S.; Buckshee, K.; Farooq, A. Reproductive functions in obese women. *Prog. Food Nutr. Sci.* **1993**, 17 (2), 89–98.
39. Goldfarb, S.; Pugh, T.D. Morphology and anatomic localization of renal microneoplasms and proximal tubule dysplasias induced by four different estrogens in the hamster. *Cancer Res.* **1990**, 50 (1), 113–119.
40. Weisz, J.; Fritz-Wolz, G.; Clawson, G.A.; Benedict, C.M.; Abendroth, C.; Creveling, C.R. Induction of nuclear catechol-O-methyltransferase by estrogens in hamster kidney: implications for estrogen-induced renal cancer. *Carcinogenesis* **1998**, 19 (7), 1307–1312.
41. Attia, N.; Tamborlane, W.V.; Heptulla, R.; Maggs, D.; Grozman, A.; Sherwin, R.S.; Caprio, S. The metabolic syndrome and insulin-like growth factor I regulation in adolescent obesity. *J. Clin. Endocrinol. Metab.* **1998**, 83 (5), 1467–1471.
42. Narayan, S.; Roy, D. Insulin-like growth factor 1 receptors are increased in estrogen-induced kidney tumors. *Cancer Res.* **1993**, 53 (Suppl. 10), 2256–2259.
43. Kasiske, B.L.; O'Donnell, M.P.; Keane, W.F. The Zucker rat model of obesity, insulin resistance, hyperlipidemia, and renal injury. *Hypertension* **1992**, 19 (Suppl. 1), I110–I115.
44. Huang, Z.S.; Chien, K.L.; Yang, C.Y.; Tsai, K.S.; Wang, C.H. Peripheral differential leukocyte counts in humans vary with hyperlipidemia, smoking, and body mass index. *Lipids* **2001**, 36 (3), 237–245.
45. Lindblad, P.; Møller, A.; Schlehofer, B. International renal-cell cancer study. V. Reproductive factors, gynecologic operations and exogenous hormones. *Int. J. Cancer* **1995**, 61, 192–198.
46. Chow, W.-H.; McLaughlin, J.K.; Mandel, J.S.; Blot, W.J.; Niwa, S.; Fraumeni, J.F., Jr. Reproductive factors and the risk of renal cell cancer among women. *Int. J. Cancer* **1995**, 60, 321–324.
47. Gago-Dominguez, M.; Castela, J.E.; Yan, J.M.; Ross, R.K.; Yu, M.C. Increased risk of renal cell carcinoma subsequent to hysterectomy. *Cancer Epidemiol. Biomark. Prev.* **1999**, 8, 999–1003.
48. Deguchi, J.; Miyamoto, M.; Okada, S. Sex hormone-dependent renal cell carcinogenesis induced by ferric nitrilotriacetate in Wistar rats. *Jpn. J. Cancer Res.* **1995**, 86, 1068–1071.
49. McLaughlin, J.K.; Chow, W.H.; Mandel, J.S.; Møller, A.; McCredie, M.; Lindblad, P.; Schlehofer, B.; Pommer, W.; Niwa, S.; Adami, H.O. International renal-cell cancer study. VIII. Role of diuretics, other anti-hypertensive medications and hypertension. *Int. J. Cancer* **1995**, 63 (2), 216–221.
50. Raynor, W.J., Jr.; Shekelle, R.B.; Ross, A.H.; Maliza, C.; Paul, O. High blood pressure and 17-year cancer mortality in the Western Electric Health Study. *Am. J. Epidemiol.* **1981**, 113 (4), 371–377.

51. Buck, C.; Donner, A. Cancer incidence in hypertensives. *Cancer* **1987**, *59* (7), 1386–1390.
52. Hole, D.J.; Hawthorne, V.M.; Isles, C.G.; McGhee, S.M.; Robertson, J.W.; Gillis, C.R.; Wapshaw, J.A.; Lever, A.F. Incidence of and mortality from cancer in hypertensive patients. *BMJ* **1993**, *306* (6878), 609–611.
53. McCredie, M.; Stewart, J.H. Risk factors for kidney cancer in New South Wales, Australia. II. Urologic disease, hypertension, obesity, and hormonal factors. *Cancer Causes Control* **1992**, *3* (4), 323–331.
54. Goldbourt, U.; Holtzman, E.; Yaari, S.; Cohen, L.; Katz, L.; Neufeld, H.N. Elevated systolic blood pressure as a predictor of long-term cancer mortality: analysis by site and histologic subtype in 10,000 middle-aged and elderly men. *J. Natl. Cancer Inst.* **1986**, *77* (1), 63–70.
55. Asal, N.R.; Geyer, J.R.; Risser, D.R.; Lee, E.T.; Kadamani, S.; Cherg, N. Risk factors in renal cell carcinoma. II. Medical history, occupation, multivariate analysis, and conclusions. *Cancer Detect. Prev.* **1988**, *13* (3–4), 263–279.
56. Chow, W.H.; McLaughlin, J.K.; Mandel, J.S.; Wacholder, S.; Niwa, S.; Fraumeni, J.F., Jr. Risk of renal cell cancer in relation to diuretics, antihypertensive drugs, and hypertension. *Cancer Epidemiol. Biomark. Prev.* **1995**, *4* (4), 327–331.
57. Møller, A.; Niwa, S.; Mehl, E.S.; Engholm, G.; McLaughlin, J.K.; Olsen, J.H. Risk factors for renal cell carcinoma in Denmark: role of medication and medical history. *Int. J. Epidemiol.* **1994**, *23* (5), 923–930.
58. McLaughlin, J.K.; Malmer, H.S.; Blot, W.J.; Weiner, J.A.; Stone, B.J.; Ericsson, J.L.; Fraumeni, J.F., Jr. Renal cell cancer among architects and allied professionals in Sweden. *Am. J. Ind. Med.* **1992**, *21* (6), 873–876.
59. Shapiro, J.A.; Williams, M.A.; Weiss, N.S.; Stergachis, A.; LaCroix, A.Z.; Barlow, W.E. Hypertension, antihypertensive medication use, and risk of renal cell carcinoma. *Am. J. Epidemiol.* **1999**, *149* (6), 521–530.
60. Federation of American Societies for Experimental Biology, L.S.R.O. *Third Report on Nutrition Monitoring in the United States*; U.S. Government Printing Office: Washington, D.C., 1995.
61. Chow, W.H.; Devesa, S.S.; Fraumeni, J.F., Jr. Epidemiology of renal cell carcinoma. Part III. Kidney cancers. In *Comprehensive Textbook of Genitourinary Oncology*; Vogelzang, N.J., Scardino, P.T., Shipley, W.U., Coffey, D.S., Miles, B.J., Eds.; Lippincott Williams & Wilkins: Philadelphia, PA, 2000.
62. Flegal, K.M.; Carroll, M.D.; Kuczmarski, R.J.; Johnson, C.L. Overweight and obesity in the United States: prevalence and trends, 1960–1994. *Int. J. Obes. Relat. Metab. Disord.* **1998**, *22* (1), 39–47.
63. Brown, C.D.; Higgins, M.; Donato, K.A.; Rohde, F.C.; Garrison, R.; Obarzanek, E.; Ernst, N.D.; Horan, M. Body mass index and the prevalence of hypertension and dyslipidemia. *Obes. Res.* **2000**, *8* (9), 605–619.
64. Chow, W.H.; McLaughlin, J.K.; Linet, M.S.; Niwa, S.; Mandel, J.S. Use of analgesics and risk of renal cell cancer. *Int. J. Cancer* **1994**, *59* (4), 467–470.
65. Matson, M.A.; Cohen, E.P. Acquired cystic kidney disease: occurrence, prevalence, and renal cancers. *Medicine (Baltimore)* **1990**, *69* (4), 217–226.
66. Levine, E.; Slusher, S.L.; Grantham, J.J.; Wetzel, L.H. Natural history of acquired renal cystic disease in dialysis patients: a prospective longitudinal CT study. *AJR Am. J. Roentgenol.* **1991**, *156* (3), 501–506.
67. Brownson, R.C. A case-control study of renal cell carcinoma in relation to occupation, smoking, and alcohol consumption. *Arch. Environ. Health* **1988**, *43* (3), 238–241.
68. Maclure, M.; Willett, W. A case-control study of diet and risk of renal adenocarcinoma. *Epidemiology* **1990**, *1* (6), 430–440.
69. McLaughlin, J.K.; Gao, Y.T.; Gao, R.N.; Zheng, W.; Ji, B.T.; Blot, W.J.; Fraumeni, J.F., Jr. Risk factors for renal-cell cancer in Shanghai, China. *Int. J. Cancer* **1992**, *52* (4), 562–565.
70. Wolk, A.; Gridley, G.; Niwa, S.; Lindblad, P.; McCredie, M.; Møller, A.; Mandel, J.S.; Wahrendorf, J.; McLaughlin, J.K.; Adami, H.O. International renal cell cancer study. VII. Role of diet. *Int. J. Cancer* **1996**, *65* (1), 67–73.
71. Lindblad, P.; Wolk, A.; Bergstrom, R.; Adami, H.-O. Diet and risk of renal cell cancer: a population-based case-control study. *Cancer Epidemiol. Biomark. Prev.* **1997**, *6*, 215–223.
72. Yuan, J.-M.; Gago-Dominguez, M.; Castella, J.E.; Hankin, J.H.; Ross, R.K.; Yu, M.C. Cruciferous vegetables in relation to renal cell carcinoma. *Int. J. Cancer* **1998**, *77*, 211–216.
73. Smith, A.H.; Shearn, V.I.; Wood, R. Asbestos and kidney cancer: the evidence supports a causal association. *Am. J. Ind. Med.* **1989**, *16* (2), 159–166.
74. Patel-Mandlik, K.J. Asbestos fibers in normal and cancerous human kidneys. *Arch. Environ. Contam. Toxicol.* **1981**, *10* (1), 47–54.
75. Huang, J.; Hisanaga, N.; Sakai, K.; Iwata, M.; Ono, Y.; Shibata, E.; Takeuchi, Y. Asbestos fibers in human pulmonary and extrapulmonary tissues. *Am. J. Ind. Med.* **1988**, *14* (3), 331–339.
76. Selikoff, I.J.; Hammond, E.C.; Seidman, H. Mortality experience of insulation workers in the United States and Canada, 1943–1976. *Ann. N.Y. Acad. Sci.* **1979**, *330*, 91–116.
77. Maclure, M. Asbestos and renal adenocarcinoma: a case-control study. *Environ. Res.* **1987**, *42* (2), 353–361.
78. Enterline, P.E.; Hartley, J.; Henderson, V. Asbestos and cancer: a cohort followed up to death. *Br. J. Ind. Med.* **1987**, *44* (6), 396–401.
79. Mandel, J.S.; McLaughlin, J.K.; Schlehofer, B.; Møller, A.; Helmer, U.; Lindblad, P.; McCredie, M.; Adami, H.O. International renal-cell cancer study. IV. Occupation. *Int. J. Cancer* **1995**, *61* (5), 601–605.
80. Puntoni, R.; Vercelli, M.; Merlo, F.; Valerio, F.; Santi, L. Mortality among shipyard workers in Genoa, Italy. *Ann. N.Y. Acad. Sci.* **1979**, *330*, 353–377.
81. Parent, M.E.; Hua, Y.; Siemiatycki, J. Occupational risk factors for renal cell carcinoma in Montreal. *Am. J. Ind. Med.* **2000**, *38* (6), 609–618.
82. Bond, G.G.; Shellenberger, R.J.; Flores, G.H.; Cook, R.R.; Fishbeck, W.A. A case-control study of renal cancer mortality at a Texas chemical plant. *Am. J. Ind. Med.* **1985**, *7* (2), 123–139.
83. Saarni, H.; Pentti, J.; Pukkala, E. Cancer at sea: a case-control study among male Finnish seafarers. *Occup. Environ. Med.* **2002**, *59* (9), 613–619.
84. Sali, D.; Boffetta, P. Kidney cancer and occupational exposure to asbestos: a meta-analysis of occupational cohort studies. *Cancer Causes Control* **2000**, *11* (1), 37–47.
85. Goodman, M.; Morgan, R.W.; Ray, R.; Malloy, C.D.; Zhao, K. Cancer in asbestos-exposed occupational cohorts: a meta-analysis. *Cancer Causes Control* **1999**, *10* (5), 453–465.
86. Hu, J.; Mao, Y.; White, K. Renal cell carcinoma and occupational exposure to chemicals in Canada. *Occup. Med. (Lond.)* **2002**, *52* (3), 157–164.
87. Partanen, T.; Heikkilä, P.; Hernberg, S.; Kauppinen, T.; Moneta, G.; Ojajarvi, A. Renal cell cancer and occupational exposure to chemical agents. *Scand. J. Work, Environ. & Health* **1991**, *17* (4), 231–239.
88. Pesch, B.; Haerting, J.; Ranft, U.; Klimpel, A.; Oelschlagel, B.; Schill, W. Occupational risk factors for renal cell carcinoma: agent-specific results from a case-control study in Germany. MURC Study Group. Multicenter urothelial and renal cancer study. *Int. J. Epidemiol.* **2000**, *29* (6), 1014–1024.
89. MacFarland, R.T.; Gandolfi, A.J.; Sipes, I.G. Extra-hepatic GSH-dependent metabolism of 1,2-dibromoethane (DBE) and 1,2-dibromo-3-chloropropane (DBCP) in the rat and mouse. *Drug Chem. Toxicol.* **1984**, *7* (3), 213–227.

90. McLaughlin, J.K.; Blot, W.J.; Mehl, E.S.; Stewart, P.A.; Venable, F.S.; Fraumeni, J.F., Jr. Petroleum-related employment and renal cell cancer. *J. Occup. Med.* **1985**, *27* (9), 672–674.
91. Lynge, E.; Anttila, A.; Hemminki, K. Organic solvents and cancer. *Cancer Causes Control* **1997**, *8* (3), 406–419.
92. Schnatter, A.R.; Katz, A.M.; Nicolich, M.J.; Theriault, G. A retrospective mortality study among Canadian petroleum marketing and distribution workers. *Environ. Health Perspect.* **1993**, *101* (Suppl. 6), 85–99.
93. Siemiatycki, J.; Dewar, R.; Nadon, L.; Gerin, M.; Richardson, L.; Wacholder, S. Associations between several sites of cancer and twelve petroleum-derived liquids. Results from a case-referent study in Montreal. *Scand. J. Work, Environ. & Health* **1987**, *13* (6), 493–504.
94. Schlehofer, B.; Heuer, C.; Blettner, M.; Niehoff, D.; Wahrendorf, J. Occupation, smoking and demographic factors, and renal cell carcinoma in Germany. *Int. J. Epidemiol.* **1995**, *24* (1), 51–57.
95. Sharpe, C.R.; Rochon, J.E.; Adam, J.M.; Suissa, S. Case-control study of hydrocarbon exposures in patients with renal cell carcinoma. *CMAJ* **1989**, *140* (11), 1309–1318.
96. McLaughlin, J.K. Renal cell cancer and exposure to gasoline: a review. *Environ. Health Perspect.* **1993**, *101* (Suppl. 6), 111–114.
97. Dement, J.M.; Hensley, L.; Gitelman, A. Carcinogenicity of gasoline: a review of epidemiological evidence. *Ann. N.Y. Acad. Sci.* **1997**, *837*, 53–76.
98. Brown, D.P.; Kaplan, S.D. Retrospective cohort mortality study of dry cleaner workers using perchloroethylene. *J. Occup. Med.* **1987**, *29* (6), 535–541.
99. Blair, A.; Decoufle, P.; Grauman, D. Causes of death among laundry and dry cleaning workers. *Am. J. Public Health* **1979**, *69* (5), 508–511.
100. Duh, R.W.; Asal, N.R. Mortality among laundry and dry cleaning workers in Oklahoma. *Am. J. Public Health* **1984**, *74* (11), 1278–1280.
101. Katz, R.M.; Jowett, D. Female laundry and dry cleaning workers in Wisconsin: a mortality analysis. *Am. J. Public Health* **1981**, *71* (3), 305–307.
102. Lowery, J.T.; Peters, J.M.; Deapen, D.; London, S.J. Renal cell carcinoma among architects. *Am. J. Ind. Med.* **1991**, *20* (1), 123–125.
103. Morgan, R.W.; Kelsh, M.A.; Zhao, K.; Heringer, S. Mortality of aerospace workers exposed to trichloroethylene. *Epidemiology* **1998**, *9* (4), 424–431.
104. Blair, A.; Hartge, P.; Stewart, P.A.; McAdams, M.; Lubin, J. Mortality and cancer incidence of aircraft maintenance workers exposed to trichloroethylene and other organic solvents and chemicals: extended follow up. *Occup. Environ. Med.* **1998**, *55* (3), 161–171.
105. Blair, A.; Stewart, P.A.; Tolbert, P.E.; Grauman, D.; Moran, F.X.; Vaught, J.; Rayner, J. Cancer and other causes of death among a cohort of dry cleaners. *Br. J. Ind. Med.* **1990**, *47* (3), 162–168.
106. McLaughlin, J.K.; Blot, W.J. A critical review of epidemiology studies of trichloroethylene and perchloroethylene and risk of renal-cell cancer. *Int. Arch. Occup. Environ. Health* **1997**, *70* (4), 222–231.
107. Blair, A.; Petralia, S.A.; Stewart, P.A. Extended mortality follow-up of a cohort of dry cleaners. *Ann. Epidemiol.* **2003**, *13* (1), 50–56.
108. Wu, C.; Schaum, J. Exposure assessment of trichloroethylene. *Environ. Health Perspect.* **2000**, *108* (Suppl. 2), 359–363.
109. Axelson, O.; Selden, A.; Andersson, K.; Hogstedt, C. Updated and expanded Swedish cohort study on trichloroethylene and cancer risk. *J. Occup. Med.* **1994**, *36* (5), 556–562.
110. Spirtas, R.; Stewart, P.A.; Lee, J.S.; Marano, D.E.; Forbes, C.D.; Grauman, D.J.; Pettigrew, H.M.; Blair, A.; Hoover, R.N.; Cohen, J.L. Retrospective cohort mortality study of workers at an aircraft maintenance facility. I. Epidemiological results. *Br. J. Ind. Med.* **1991**, *48* (8), 515–530.
111. Anttila, A.; Pukkala, E.; Sallmen, M.; Hernberg, S.; Hemminki, K. Cancer incidence among Finnish workers exposed to halogenated hydrocarbons. *J. Occup. Environ. Med.* **1995**, *37* (7), 797–806.
112. Garabrant, D.H.; Held, J.; Langholz, B.; Bernstein, L. Mortality of aircraft manufacturing workers in southern California. *Am. J. Ind. Med.* **1988**, *13* (6), 683–693.
113. Henschler, D.; Vamvakas, S.; Lammert, M.; Dekant, W.; Kraus, B.; Thomas, B.; Ulm, K. Increased incidence of renal cell tumors in a cohort of cardboard workers exposed to trichloroethene. *Arch. Toxicol.* **1995**, *69* (5), 291–299.
114. Vamvakas, S.; Bruning, T.; Thomasson, B.; Lammert, M.; Baumuller, A.; Bolt, H.M.; Dekant, W.; Birner, G.; Henschler, D.; Ulm, K. Renal cell cancer correlated with occupational exposure to trichloroethene. *J. Cancer Res. Clin. Oncol.* **1998**, *124* (7), 374–382.
115. Mandel, J.S. Renal cell cancer correlated with occupational exposure to trichloroethylene. *J. Cancer Res. Clin. Oncol.* **2001**, *127* (4), 265–268.
116. Bruning, T.; Pesch, B.; Wiesenhuber, B.; Rabstein, S.; Lammert, M.; Baumuller, A.; Bolt, H.M. Renal cell cancer risk and occupational exposure to trichloroethylene: results of a consecutive case-control study in Arnsberg, Germany. *Am. J. Ind. Med.* **2003**, *43*, 274–285.
117. Dosemeci, M.; Cocco, P.; Chow, W.H. Gender differences in risk of renal cell carcinoma and occupational exposures to chlorinated aliphatic hydrocarbons. *Am. J. Ind. Med.* **1999**, *36* (1), 54–59.
118. Gomez, M.R.; Cocco, P.; Dosemeci, M.; Stewart, P.A. Occupational exposure to chlorinated aliphatic hydrocarbons: job exposure matrix. *Am. J. Ind. Med.* **1994**, *26* (2), 171–183.
119. Ruder, A.M.; Ward, E.M.; Brown, D.P. Cancer mortality in female and male dry-cleaning workers. *J. Occup. Med.* **1994**, *36* (8), 867–874.
120. Hansen, J.; Raaschou-Nielsen, O.; Christensen, J.M.; Johansen, I.; McLaughlin, J.K.; Lipworth, L.; Blot, W.J.; Olsen, J.H. Cancer incidence among Danish workers exposed to trichloroethylene. *J. Occup. Environ. Med.* **2001**, *43* (2), 133–139.
121. Brauch, H.; Weirich, G.; Hornauer, M.A.; Storkel, S.; Wohl, T.; Bruning, T. Trichloroethylene exposure and specific somatic mutations in patients with renal cell carcinoma. *J. Natl. Cancer Inst.* **1999**, *91* (10), 854–861.
122. Bruning, T.; Bolt, H.M. Renal toxicity and carcinogenicity of trichloroethylene: key results, mechanisms, and controversies. *Crit. Rev. Toxicol.* **2000**, *30* (3), 253–285.
123. Dekant, W. Chemical-induced nephrotoxicity mediated by glutathione S-conjugate formation. *Toxicol. Lett.* **2001**, *124* (1–3), 21–36.
124. Boffetta, P.; Dosemeci, M.; Gridley, G.; Bath, H.; Moradi, T.; Silverman, D. Occupational exposure to diesel engine emissions and risk of cancer in Swedish men and women. *Cancer Causes Control* **2001**, *12* (4), 365–374.
125. Soll-Johanning, H.; Bach, E.; Olsen, J.H.; Tuchsien, F. Cancer incidence in urban bus drivers and tramway employees: a retrospective cohort study. *Occup. Environ. Med.* **1998**, *55* (9), 594–598.
126. Redmond, C.K.; Ciocco, A.; Lloyd, J.W.; Rush, H.W. Long-term mortality study of steelworkers. VI. Mortality from malignant neoplasms among coke oven workers. *J. Occup. Med.* **1972**, *14* (8), 621–629.
127. Redmond, C.K. Cancer mortality among coke oven workers. *Environ. Health Perspect.* **1983**, *52*, 67–73.
128. Costantino, J.P.; Redmond, C.K.; Bearden, A. Occupationally related cancer risk among coke oven workers: 30 years of follow-up. *J. Occup. Environ. Med.* **1995**, *37* (5), 597–604.
129. Bertazzi, P.A.; Pesatori, A.C.; Zocchetti, C.; Latocca, R. Mortality study of cancer risk among oil refinery workers. *Int. Arch. Occup. Environ. Health* **1989**, *61* (4), 261–270.
130. Savitz, D.A.; Moure, R. Cancer risk among oil refinery workers. A review of epidemiologic studies. *J. Occup. Med.* **1984**, *26* (9), 662–670.

131. Wong, O.; Harris, F.; Smith, T.J. Health effects of gasoline exposure. II. Mortality patterns of distribution workers in the United States. *Environ. Health Perspect.* **1993**, *101* (Suppl. 6), 63–76.
132. Gamble, J.F.; Lewis, R.J.; Jorgensen, G. Mortality among three refinery/petrochemical plant cohorts. II. Retirees. *J. Occup. Environ. Med.* **2000**, *42* (7), 730–736.
133. Wong, O.; Raabe, G.K. A critical review of cancer epidemiology in the petroleum industry, with a meta-analysis of a combined database of more than 350,000 workers. *Regul. Toxicol. Pharmacol.* **2000**, *32* (1), 78–98.
134. Pukkala, E. Cancer incidence among Finnish oil refinery workers, 1971–1994. *J. Occup. Environ. Med.* **1998**, *40* (8), 675–679.
135. Poole, C.; Dreyer, N.A.; Satterfield, M.H.; Levin, L.; Rothman, K.J. Kidney cancer and hydrocarbon exposures among petroleum refinery workers. *Environ. Health Perspect.* **1993**, *101* (Suppl. 6), 53–62.
136. Rushton, L. A 39-year follow-up of the U.K. oil refinery and distribution center studies: results for kidney cancer and leukemia. *Environ. Health Perspect.* **1993**, *101* (Suppl. 6), 77–84.
137. Thomas, T.L.; Waxweiler, R.J.; Moure-Eraso, R.; Itaya, S.; Fraumeni, J.F., Jr. Mortality patterns among workers in three Texas oil refineries. *J. Occup. Med.* **1982**, *24* (2), 135–141.
138. Ronneberg, A.; Langmark, F. Epidemiologic evidence of cancer in aluminum reduction plant workers. *Am. J. Ind. Med.* **1992**, *22* (4), 573–590.
139. Ronneberg, A.; Andersen, A. Mortality and cancer morbidity in workers from an aluminium smelter with prebaked carbon anodes. Part II. Cancer morbidity. *Occup. Environ. Med.* **1995**, *52* (4), 250–254.
140. Rockette, H.E.; Arena, V.C. Mortality studies of aluminum reduction plant workers: potroom and carbon department. *J. Occup. Med.* **1983**, *25* (7), 549–557.
141. Boffetta, P.; Jourenkova, N.; Gustavsson, P. Cancer risk from occupational and environmental exposure to polycyclic aromatic hydrocarbons. *Cancer Causes Control* **1997**, *8* (3), 444–472.
142. Mason, T.J. Geographic patterns of cancer risk: a means for identifying possible occupational factors. *Ann. N.Y. Acad. Sci.* **1976**, *271*, 370–376.
143. Lloyd, J.W.; Decoufle, P.; Salvin, L.G. Unusual mortality experience of printing pressmen. *J. Occup. Med.* **1977**, *19* (8), 543–550.
144. Thomas, T.L.; Decoufle, P.; Moure-Eraso, R. Mortality among workers employed in petroleum refining and petrochemical plants. *J. Occup. Med.* **1980**, *22* (2), 97–103.
145. Sinks, T.; Lushniak, B.; Haussler, B.J.; Sniezek, J.; Deng, J.F.; Roper, P.; Dill, P.; Coates, R. Renal cell cancer among paperboard printing workers. *Epidemiology* **1992**, *3* (6), 483–489.
146. Paganini-Hill, A.; Glazer, E.; Henderson, B.E.; Ross, R.K. Cause-specific mortality among newspaper web pressmen. *J. Occup. Med.* **1980**, *22* (8), 542–544.
147. Greene, M.H.; Hoover, R.N.; Eck, R.L.; Fraumeni, J.F., Jr. Cancer mortality among printing plant workers. *Environ. Res.* **1979**, *20* (1), 66–73.
148. Auperin, A.; Benhamou, S.; Ory-Paoletti, C.; Flamant, R. Occupational risk factors for renal cell carcinoma: a case-control study. *Occup. Environ. Med.* **1994**, *51* (6), 426–428.
149. Malke, H.R.; Malke, B.K.; McLaughlin, J.K.; Blot, W.J. Kidney cancer among leather workers. *Lancet* **1984**, *1* (8367), 56.
150. Kolonel, L.N. Association of cadmium with renal cancer. *Cancer* **1976**, *37* (4), 1782–1787.
151. Elinder, C.G.; Kjellstrom, T.; Hogstedt, C.; Andersson, K.; Spang, G. Cancer mortality of cadmium workers. *Br. J. Ind. Med.* **1985**, *42* (10), 651–655.
152. Thun, M.J.; Schnorr, T.M.; Smith, A.B.; Halperin, W.E.; Lemen, R.A. Mortality among a cohort of U.S. cadmium production workers—an update. *J. Natl. Cancer Inst.* **1985**, *74* (2), 325–333.
153. Lemen, R.A.; Lee, J.S.; Wagoner, J.K.; Blejer, H.P. Cancer mortality among cadmium production workers. *Ann. N.Y. Acad. Sci.* **1976**, *271*, 273–279.
154. Kjellstrom, T.; Friberg, L.; Rahnster, B. Mortality and cancer morbidity among cadmium-exposed workers. *Environ. Health Perspect.* **1979**, *28*, 199–204.
155. Silbergeld, E.K.; Waalkes, M.; Rice, J.M. Lead as a carcinogen: experimental evidence and mechanisms of action. *Am. J. Ind. Med.* **2000**, *38* (3), 316–323.
156. Fowler, B.A.; Kahng, M.W.; Smith, D.R. Role of lead-binding proteins in renal cancer. *Environ. Health Perspect.* **1994**, *102* (Suppl. 3), 115–116.
157. Steenland, K.; Boffetta, P. Lead and cancer in humans: where are we now? *Am. J. Ind. Med.* **2000**, *38* (3), 295–299.
158. Steenland, K.; Selevan, S.; Landrigan, P. The mortality of lead smelter workers: an update. *Am. J. Public Health* **1992**, *82* (12), 1641–1644.
159. Selevan, S.G.; Landrigan, P.J.; Stern, F.B.; Jones, J.H. Mortality of lead smelter workers. *Am. J. Epidemiol.* **1985**, *122* (4), 673–683.
160. Cocco, P.; Hua, F.; Boffetta, P.; Carta, P.; Flore, C.; Flore, V.; Onnis, A.; Picchiri, G.F.; Colin, D. Mortality of Italian lead smelter workers. *Scand. J. Work, Environ. & Health* **1997**, *23* (1), 15–23.
161. McLaughlin, J.K.; Malke, H.S.; Stone, B.J.; Weiner, J.A.; Malke, B.K.; Ericsson, J.L.; Blot, W.J.; Fraumeni, J.F., Jr. Occupational risks for renal cancer in Sweden. *Br. J. Ind. Med.* **1987**, *44* (2), 119–123.
162. Simonato, L.; Fletcher, A.C.; Andersen, A.; Anderson, K.; Becker, N.; Chang-Claude, J.; Ferro, G.; Gerin, M.; Gray, C.N.; Hansen, K.S. A historical prospective study of European stainless steel, mild steel, and shipyard welders. *Br. J. Ind. Med.* **1991**, *48* (3), 145–154.
163. Stern, F.B.; Sweeney, M.H.; Ward, E. Proportionate mortality among unionized construction ironworkers. *Am. J. Ind. Med.* **1997**, *31* (2), 176–187.
164. Hemminki, K.; Jiang, Y.; Ma, X.; Yang, K.; Egevad, L.; Lindblad, P. Molecular epidemiology of VHL gene mutations in renal cell carcinoma patients: relation to dietary and other factors. *Carcinogenesis* **2002**, *23* (5), 809–815.
165. Cooper, W.C.; Wong, O.; Kheifets, L. Mortality among employees of lead battery plants and lead-producing plants, 1947–1980. *Scand. J. Work, Environ. & Health* **1985**, *11* (5), 331–345.
166. Sankila, R.; Karjalainen, S.; Laara, E.; Pukkala, E.; Teppo, L. Cancer risk among health care personnel in Finland, 1971–1980. *Scand. J. Work, Environ. & Health* **1990**, *16* (4), 252–257.
167. Lerer, T.J.; Redmond, C.K.; Breslin, P.P.; Salvin, L.; Rush, H.W. Long-term mortality study of steelworkers. VII. Mortality patterns among crane operators. *J. Occup. Med.* **1974**, *16* (9), 608–614.
168. Andersen, A.; Barlow, L.; Engeland, A.; Kjaerheim, K.; Lynge, E.; Pukkala, E. Work-related cancer in the Nordic countries. *Scand. J. Work, Environ. & Health* **1999**, *25* (Suppl. 2), 1–116.
169. Mattos, I.E.; Sauaia, N.; Menezes, P.R. A cancer mortality pattern in Brazilian electrical workers. *Cad. Saude Publica* **2002**, *18* (1), 221–233.
170. Shalat, S.L.; True, L.D.; Fleming, L.E.; Pace, P.E. Kidney cancer in utility workers exposed to polychlorinated biphenyls (PCBs). *Br. J. Ind. Med.* **1989**, *46* (11), 823–824.
171. Milham, S., Jr. Cancer mortality pattern associated with exposure to metals. *Ann. N.Y. Acad. Sci.* **1976**, *271*, 243–249.
172. Delahunt, B.; Bethwaite, P.B.; Nacey, J.N. Occupational risk for renal cell carcinoma. A case-control study based on the New Zealand Cancer Registry. *Br. J. Urol.* **1995**, *75* (5), 578–582.
173. Guidotti, T.L. Occupational mortality among firefighters: assessing the association. *J. Occup. Environ. Med.* **1995**, *37* (12), 1348–1356.
174. Nicholas, J.S.; Lackland, D.T.; Dosemeci, M.; Mohr, L.C., Jr.; Dunbar, J.B.; Grosche, B.; Hoel, D.G. Mortality among US commercial pilots and navigators. *J. Occup. Environ. Med.* **1998**, *40* (11), 980–985.

175. Band, P.R.; Le, N.D.; Fang, R.; Threlfall, W.J.; Astrakianakis, G.; Anderson, J.T.; Keefe, A.; Krewski, D. Cohort mortality study of pulp and paper mill workers in British Columbia, Canada. *Am. J. Epidemiol.* **1997**, *146* (2), 186–194.
176. Mattioli, S.; Truffelli, D.; Baldasseroni, A.; Risi, A.; Marchesini, B.; Giacomini, C.; Bacchini, P.; Violante, F.S.; Buiatti, E. Occupational risk factors for renal cell cancer: a case-control study in northern Italy. *J. Occup. Environ. Med.* **2002**, *44* (11), 1028–1036.
177. Bruning, T.; Chronz, C.; Thier, R.; Havelka, J.; Ko, Y.; Bolt, H.M. Occurrence of urinary tract tumors in miners highly exposed to dinitrotoluene. *J. Occup. Environ. Med.* **1999**, *41* (3), 144–149.
178. Enterline, P.E.; Day, R.; Marsh, G.M. Cancers related to exposure to arsenic at a copper smelter. *Occup. Environ. Med.* **1995**, *52* (1), 28–32.
179. Cantor, K.P.; Sontag, J.M.; Heid, M.F. Patterns of mortality among plumbers and pipefitters. *Am. J. Ind. Med.* **1986**, *10* (1), 73–89.
180. Miller, B.A.; Blair, A.; Reed, E.J. Extended mortality follow-up among men and women in a U.S. furniture workers union. *Am. J. Ind. Med.* **1994**, *25* (4), 537–549.
181. Buzio, L.; Tondel, M.; De Palma, G.; Buzio, C.; Franchini, I.; Mutti, A.; Axelson, O. Occupational risk factors for renal cell cancer. An Italian case-control study. *Med. Lav.* **2002**, *93* (4), 303–309.
182. Cantor, K.P.; Hoover, R.; Mason, T.J.; McCabe, L.J. Associations of cancer mortality with halomethanes in drinking water. *J. Natl. Cancer Inst.* **1978**, *61* (4), 979–985.
183. Wilkins, J.R., III; Comstock, G.W. Source of drinking water at home and site-specific cancer incidence in Washington County, Maryland. *Am. J. Epidemiol.* **1981**, *114* (2), 178–190.
184. Yang, C.Y.; Chiu, H.F.; Cheng, M.F.; Tsai, S.S. Chlorination of drinking water and cancer mortality in Taiwan. *Environ. Res.* **1998**, *78* (1), 1–6.
185. Koivusalo, M.; Hakulinen, T.; Vartiainen, T.; Pukkala, E.; Jaakkola, J.J.; Tuomisto, J. Drinking water mutagenicity and urinary tract cancers: a population-based case-control study in Finland. *Am. J. Epidemiol.* **1998**, *148* (7), 704–712.
186. Chen, C.J.; Chuang, Y.C.; Lin, T.M.; Wu, H.Y. Malignant neoplasms among residents of a blackfoot disease-endemic area in Taiwan: high-arsenic artesian well water and cancers. *Cancer Res.* **1985**, *45* (11 Pt 2), 5895–5899.
187. Wu, M.M.; Kuo, T.L.; Hwang, Y.H.; Chen, C.J. Dose-response relation between arsenic concentration in well water and mortality from cancers and vascular diseases. *Am. J. Epidemiol.* **1989**, *130* (6), 1123–1132.
188. Chen, C.J.; Wang, C.J. Ecological correlation between arsenic level in well water and age-adjusted mortality from malignant neoplasms. *Cancer Res.* **1990**, *50* (17), 5470–5474.
189. Hoppenhay-Rich, C.; Biggs, M.L.; Smith, A.H. Lung and kidney cancer mortality associated with arsenic in drinking water in Cordoba, Argentina. *Int. J. Epidemiol.* **1998**, *27* (4), 561–569.
190. Chiou, H.Y.; Chiou, S.T.; Hsu, Y.H.; Chou, Y.L.; Tseng, C.H.; Wei, M.L.; Chen, C.J. Incidence of transitional cell carcinoma and arsenic in drinking water: a follow-up study of 8,102 residents in an arseniasis-endemic area in northeastern Taiwan. *Am. J. Epidemiol.* **2001**, *153* (5), 411–418.
191. Kurtio, P.; Pukkala, E.; Kahelin, H.; Auvinen, A.; Pekkanen, J. Arsenic concentrations in well water and risk of bladder and kidney cancer in Finland. *Environ. Health Perspect.* **1999**, *107* (9), 705–710.
192. Howe, H.L.; Wolfgang, P.E.; Burnett, W.S.; Nasca, P.C.; Youngblood, L. Cancer incidence following exposure to drinking water with asbestos leachate. *Public Health Rep.* **1989**, *104* (3), 251–256.
193. Sadler, T.D.; Rom, W.N.; Lyon, J.L.; Mason, J.O. The use of asbestos-cement pipe for public water supply and the incidence of cancer in selected communities in Utah. *J. Commun. Health* **1984**, *9* (4), 285–293.
194. Kanarek, M.S.; Conforti, P.M.; Jackson, L.A.; Cooper, R.C.; Murchio, J.C. Asbestos in drinking water and cancer incidence in the San Francisco Bay area. *Am. J. Epidemiol.* **1980**, *112* (1), 54–72.
195. Graham, S.; Blanchet, M.; Rohrer, T. Cancer in asbestos-mining and other areas of Quebec. *J. Natl. Cancer Inst.* **1977**, *59* (4), 1139–1145.
196. Cantor, K.P. Invited commentary: arsenic and cancer of the urinary tract. *Am. J. Epidemiol.* **2001**, *153* (5), 422–423.
197. Phillips, J.L.; Pavlovich, C.P.; Walther, M.; Ried, T.; Linehan, W.M. The genetic basis of renal epithelial tumors: advances in research and its impact on prognosis and therapy. *Curr. Opin. Urol.* **2001**, *11* (5), 463–469.
198. Latif, F.; Tory, K.; Gnarr, J.; Yao, M.; Duh, F.M.; Orcutt, M.L.; Stackhouse, T.; Kuzmin, I.; Modi, W.; Geil, L. Identification of the von Hippel-Lindau disease tumor suppressor gene. *Science* **1993**, *260* (5112), 1317–1320.
199. Pause, A.; Peterson, B.; Schaffar, G.; Stearman, R.; Klausner, R.D. Studying interactions of four proteins in the yeast two-hybrid system: structural resemblance of the pVHL/elongin BC/hCUL-2 complex with the ubiquitin ligase complex SKP1/cullin/F-box protein. *Proc. Natl. Acad. Sci. U. S. A.* **1999**, *96* (17), 9533–9538.
200. Hughson, M.D.; He, Z.; Liu, S.; Coleman, J.; Shingleton, W.B. Expression of HIF-1 and ubiquitin in conventional renal cell carcinoma: relationship to mutations of the von Hippel-Lindau tumor suppressor gene. *Cancer Genet. Cytogenet.* **2003**, *143*, 145–153.
201. Pugh, C.W.; Ratcliffe, J. The von Hippel-Lindau tumor suppressor, hypoxia-inducible factor-1 (HIF-1) degradation, and cancer pathogenesis. *Semin. Cancer Biol.* **2003**, *13*, 83–89.
202. Sano, T.; Horiguchi, H. Von Hippel-Lindau disease. *Microsc. Res. Tech.* **2003**, *60* (2), 159–164.
203. Hergovich, A.; Lisztwan, J.; Barry, R.; Ballschmieter, P.; Krek, W. Regulation of microtubule stability by the von Hippel-Lindau tumour suppressor protein pVHL. *Nat. Cell Biol.* **2003**, *5* (1), 64–70.
204. Gemmill, R.M.; West, J.D.; Boldog, F.; Tanaka, N.; Robinson, L.J.; Smith, D.I.; Li, F.; Drabkin, H.A. The hereditary renal cell carcinoma 3:8 translocation fuses FHIT to a patched-related gene, TRC8. *Proc. Natl. Acad. Sci. U. S. A.* **1998**, *95* (16), 9572–9577.
205. Gallou, C.; Longueaux, S.; Delomenie, C.; Mejean, A.; Martin, N.; Martinet, S.; Palais, G.; Bouvier, R.; Droz, D.; Krishnamoorthy, R.; Junien, C.; Beroud, C.; Dupret, J.M. Association of GSTT1 non-null and NAT1 slow/rapid genotypes with von Hippel-Lindau tumour suppressor gene transversions in sporadic renal cell carcinoma. *Pharmacogenetics* **2001**, *11* (6), 521–535.
206. Ma, X.; Yang, K.; Lindblad, P.; Egevad, L.; Hemminki, K. VHL gene alterations in renal cell carcinoma patients: novel hotspot or founder mutations and linkage disequilibrium. *Oncogene* **2001**, *20* (38), 5393–5400.
207. Foster, K.; Prowse, A.; van den, B.A.; Fleming, S.; Hulsbeek, M.M.; Crossey, P.A.; Richards, F.M.; Cairns, P.; Affara, N.A.; Ferguson-Smith, M.A. Somatic mutations of the von Hippel-Lindau disease tumour suppressor gene in non-familial clear cell renal carcinoma. *Hum. Mol. Genet.* **1994**, *3* (12), 2169–2173.
208. Clifford, S.C.; Prowse, A.H.; Affara, N.A.; Buys, C.H.; Maher, E.R. Inactivation of the von Hippel-Lindau (VHL) tumour suppressor gene and allelic losses at chromosome arm 3p in primary renal cell carcinoma: evidence for a VHL-independent pathway in clear cell renal tumorigenesis. *Genes Chromosomes Cancer* **1998**, *22* (3), 200–209.
209. Bruning, T.; Weirich, G.; Hornauer, M.A.; Hofler, H.; Brauch, H. Renal cell carcinomas in trichloroethene (TRI) exposed persons are associated with somatic mutations in the von Hippel-Lindau (VHL) tumour suppressor gene. *Arch. Toxicol.* **1997**, *71* (5), 332–335.
210. Brauch, H.; Weirich, G.; Brieger, J.; Glavac, D.; Rodl, H.; Eichinger, M.; Feurer, M.; Weidt, E.; Puranakanittha, C.; Neuhaus, C.; Pomer, S.;

- Brenner, W.; Schirmacher, P.; Storkel, S.; Rotter, M.; Masera, A.; Gugeler, N.; Decker, H.J. VHL alterations in human clear cell renal cell carcinoma: association with advanced tumor stage and a novel hot spot mutation. *Cancer Res.* **2000**, *60* (7), 1942–1948.
211. Fischer, J.; Palmedo, G.; von Knobloch, R.; Bugert, P.; Prayer-Galetti, T.; Pagano, F.; Kovacs, G. Duplication and overexpression of the mutant allele of the MET proto-oncogene in multiple hereditary papillary renal cell tumours. *Oncogene* **1998**, *17* (6), 733–739.
 212. Johnson, M.; Koukoulis, G.; Matsumoto, K.; Nakamura, T.; Iyer, A. Hepatocyte growth factor induces proliferation and morphogenesis in nonparenchymal epithelial liver cells. *Hepatology* **1993**, *17* (6), 1052–1061.
 213. Tomlinson, I.P.; Alam, N.A.; Rowan, A.J.; Barclay, E.; Jaeger, E.E.; Kelsell, D.; Leigh, I.; Gorman, P.; Lamlum, H.; Rahman, S.; Roylance, R.R.; Olpin, S.; Bevan, S.; Barker, K.; Hearle, N.; Houlston, R.S.; Kiuru, M.; Lehtonen, R.; Karhu, A.; Vilkki, S.; Laiho, P.; Eklund, C.; Vierimaa, O.; Aittomaki, K.; Hietala, M.; Sistonen, P.; Paetau, A.; Salovaara, R.; Herva, R.; Launonen, V.; Aaltonen, L.A. Germline mutations in FH predispose to dominantly inherited uterine fibroids, skin leiomyomata and papillary renal cell cancer. *Nat. Genet.* **2002**, *30* (4), 406–410.
 214. Choyke, P.L.; Glenn, G.M.; Walther, M.M.; Zbar, B.; Linehan, W.M. Hereditary renal cancers. *Radiology* **2003**, *226* (1), 33–46.
 215. Nickerson, M.L.; Warren, M.B.; Toro, J.R.; Matrosova, V.; Glenn, G.; Turner, M.L.; Duray, P.; Merino, M.; Choyke, P.; Pavlovich, C.P.; Sharma, N.; Walther, M.; Munroe, D.; Hill, R.; Maher, E.; Greenberg, C.; Lerman, M.I.; Linehan, W.M.; Zbar, B.; Schmidt, L.S. Mutations in a novel gene lead to kidney tumors, lung wall defects, and benign tumors of the hair follicle in patients with the Birt-Hogg-Dube syndrome. *Cancer Cell* **2002**, *2* (2), 157–164.
 216. Schulz, W.A.; Krummeck, A.; Rosinger, I.; Eickelmann, P.; Neuhaus, C.; Ebert, T.; Schmitz-Drager, B.J.; Sies, H. Increased frequency of a null-allele for NAD(P)H: quinone oxidoreductase in patients with urological malignancies. *Pharmacogenetics* **1997**, *7* (3), 235–239.
 217. Sweeney, C.; Farrow, D.C.; Schwartz, S.M.; Eaton, D.L.; Checkoway, H.; Vaughan, T.L. Glutathione S-transferase M1, T1, and P1 polymorphisms as risk factors for renal cell carcinoma: a case-control study. *Cancer Epidemiol. Biomark. Prev.* **2000**, *9* (4), 449–454.
 218. Bruning, T.; Lammert, M.; Kempkes, M.; Thier, R.; Golka, K.; Bolt, H.M. Influence of polymorphisms of GSTM1 and GSTT1 for risk of renal cell cancer in workers with long-term high occupational exposure to trichloroethene. *Arch. Toxicol.* **1997**, *71* (9), 596–599.
 219. Semenza, J.C.; Ziogas, A.; Largent, J.; Peel, D.; Anton-Culver, H. Gene-environment interactions in renal cell carcinoma. *Am. J. Epidemiol.* **2001**, *153* (9), 851–859.
 220. Longueux, S.; Delomenie, C.; Gallou, C.; Mejean, A.; Vincent-Viry, M.; Bouvier, R.; Droz, D.; Krishnamoorthy, R.; Galteau, M.M.; Junien, C.; Beroud, C.; Dupret, J.M. Candidate genetic modifiers of individual susceptibility to renal cell carcinoma: a study of polymorphic human xenobiotic-metabolizing enzymes. *Cancer Res.* **1999**, *59* (12), 2903–2908.
 221. Gago-Dominguez, M.; Castela, J.E.; Yuan, J.-M.; Ross, R.K.; Yu, M.C. Lipid peroxidation: a novel and unifying concept of the etiology of renal cell carcinoma (United States). *Cancer Causes Control* **2002**, *13*, 287–293.
 222. Mellemgaard, A.; McLaughlin, J.K.; Overvad, K.; Olsen, J.H. Dietary risk factors for renal cell carcinoma in Denmark. *Eur. J. Cancer* **1996**, *32A* (4), 673–682.
 223. Bergstrom, A.; Terry, P.; Lindblad, P.; Lichtenstein, P.; Ahlbom, A.; Feychting, M.; Wolk, A. Physical activity and risk of renal cell cancer. *Int. J. Cancer* **2001**, *92* (1), 155–157.
 224. Lindblad, P.; Wolk, A.; Bergstrom, R.; Persson, I.; Adami, H.O. The role of obesity and weight fluctuations in the etiology of renal cell cancer: a population-based case-control study. *Cancer Epidemiol. Biomark. Prev.* **1994**, *3* (8), 631–639.
 225. Bianchi, G.D.; Cerhan, J.R.; Parker, A.S.; Putnam, S.D.; See, W.A.; Lynch, C.F.; Cantor, K.P. Tea consumption and risk of bladder and kidney cancers in a population-based case-control study. *Am. J. Epidemiol.* **2000**, *151* (4), 377–383.
 226. Chiu, B.C.; Lynch, C.F.; Cerhan, J.R.; Cantor, K.P. Cigarette smoking and risk of bladder, pancreas, kidney, and colorectal cancers in Iowa. *Ann. Epidemiol.* **2001**, *11* (1), 28–37.
 227. Parker, A.S.; Cerhan, J.R.; Lynch, C.F.; Ershow, A.G.; Cantor, K.P. Gender, alcohol consumption, and renal cell carcinoma. *Am. J. Epidemiol.* **2002**, *155* (5), 455–462.
 228. Gago-Dominguez, M.; Yuan, J.M.; Castela, J.E.; Ross, R.K.; Yu, M.C. Family history and risk of renal cell carcinoma. *Cancer Epidemiol. Biomark. Prev.* **2001**, *10* (9), 1001–1004.
 229. Yuan, J.M.; Castela, J.E.; Gago-Dominguez, M.; Ross, R.K.; Yu, M.C. Hypertension, obesity and their medications in relation to renal cell carcinoma. *Br. J. Cancer* **1998**, *77* (9), 1508–1513.
 230. Yuan, J.M.; Gago-Dominguez, M.; Castela, J.E.; Hankin, J.H.; Ross, R.K.; Yu, M.C. Cruciferous vegetables in relation to renal cell carcinoma. *Int. J. Cancer* **1998**, *77* (2), 211–216.
 231. Yuan, J.M.; Castela, J.E.; Gago-Dominguez, M.; Yu, M.C.; Ross, R.K. Tobacco use in relation to renal cell carcinoma. *Cancer Epidemiol. Biomark. Prev.* **1998**, *7* (5), 429–433.
 232. Mellemgaard, A.; Lindblad, P.; Schlehofer, B.; Bergstrom, R.; Mandel, J.S.; McCredie, M.; McLaughlin, J.K.; Niwa, S.; Odaka, N.; Pommer, W. International renal-cell cancer study. III. Role of weight, height, physical activity, and use of amphetamines. *Int. J. Cancer* **1995**, *60* (3), 350–354.
 233. Lindblad, P.; Mellemgaard, A.; Schlehofer, B.; Adami, H.O.; McCredie, M.; McLaughlin, J.K.; Mandel, J.S. International renal-cell cancer study. V. Reproductive factors, gynecologic operations and exogenous hormones. *Int. J. Cancer* **1995**, *61* (2), 192–198.
 234. Schlehofer, B.; Pommer, W.; Mellemgaard, A.; Stewart, J.H.; McCredie, M.; Niwa, S.; Lindblad, P.; Mandel, J.S.; McLaughlin, J.K.; Wahrendorf, J. International renal-cell cancer study. VI. The role of medical and family history. *Int. J. Cancer* **1996**, *6* (6), 723–726.